

## Field Treatment

1. Basic airway/spinal immobilization prn/control major bleeding prn
2. Oxygen prn
3. Hyperventilation prn  
①

4. Advanced airway prn

**Note:** ②

### PERFUSING

5. Venous access prn

**Note:** ③

6. Reverse Trendelenburg position prn
7. Reassess for potential deterioration

### POOR PERFUSION

5. Cardiac monitor/document rhythm and attach EKG strip
6. Shock position prn
7. Venous access
8. Perform blood glucose test prn

**Note:** ③ ④

9. Fluid resuscitate

## Special Considerations

- ① Hyperventilation if signs or symptoms of elevated ICP.
  - ✓ Altered LOC
  - ✓ posturing
  - ✓ bradycardia
  - ✓ hypertension
  - ✓ projectile vomiting
  - ✓ pupil dilation
  - ✓ seizure
- ② DO NOT delay transport for treatment.
- ③ Assess for internal injuries and treat by appropriate guideline.
- ④ Drugs for altered LOC with specific history:
  - ✓ Diabetic with hypoglycemia – **dextrose 50% 50 ml IVP**
  - ✓ Narcotic overdose – naloxone (Narcan®) 0.8-2 mg IVP titrated to adequate respiratory rate and tidal volume or 2mg IN/IM.